

AGE CONSTRAINTS FOR AN M31 GLOBULAR CLUSTER FROM MAIN SEQUENCE PHOTOMETRY¹

THOMAS M. BROWN², HENRY C. FERGUSON², ED SMITH², RANDY A. KIMBLE³, ALLEN V. SWEIGART³,
 ALVIO RENZINI⁴, R. MICHAEL RICH⁵, & DON A. VANDENBERG⁶

To appear in The Astrophysical Journal Letters

ABSTRACT

We present a color-magnitude diagram (CMD) of the globular cluster SKHB-312 in the Andromeda galaxy (M31), obtained with the Advanced Camera for Surveys on the Hubble Space Telescope. The cluster was included in deep observations taken to measure the star formation history of the M31 halo. Overcoming a very crowded field, our photometry of SKHB-312 reaches $m_V \approx 30.5$ mag, more than 1 mag below the main sequence turnoff. These are the first observations to allow a direct age estimate from the turnoff in an old M31 cluster. We analyze its CMD and luminosity function using a finely-spaced grid of isochrones that have been calibrated using observations of Galactic clusters taken with the same camera and filters. The luminosity difference between the subgiant and horizontal branches is ≈ 0.2 mag smaller in SKHB-312 than in the Galactic clusters 47 Tuc and NGC 5927, implying SKHB-312 is 2–3 Gyr younger. A quantitative comparison to isochrones yields an age of $10_{-1}^{+2.5}$ Gyr.

Subject headings: galaxies: star clusters – galaxies: stellar content – galaxies: halos – galaxies: individual (M31)

1. INTRODUCTION

For more than 50 years (e.g., Sandage 1953), main sequence photometry has been the most direct method for determining the ages of both star clusters and populations with more complex star formation histories. Current age estimates for globular clusters in the Galactic halo generally run from 11–13.5 Gyr (e.g., Vandenberg 2000). Although main sequence photometry has enabled the dating of globular clusters in the satellites of the Milky Way (e.g., Rich et al. 2001), no globular clusters have been imaged down to the main sequence in another spiral galaxy. Instead, extragalactic globular cluster ages are inferred from composite colors and/or spectroscopy. Recent evidence suggests that Andromeda (M31) may have more young globular clusters than the Galaxy, implying that M31 may have obtained more of its clusters from cannibalized dwarf galaxies (Barmby & Huchra 2000; Burstein et al. 2004).

We obtained extremely deep images of M31 with the Advanced Camera for Surveys (ACS) on the Hubble Space Telescope (HST) in order to measure the star formation history in its halo (Brown et al. 2003). Our field, 51' from the nucleus on the southeast minor axis, was chosen to include the M31 globular cluster SKHB-312 (Sargent et al. 1977). Spectroscopy (Huchra et al. 1991) and red giant branch (RGB) photometry (Holland et al. 1997) yield respective metallicity estimates of $[\text{Fe}/\text{H}] = -0.7$ and $[\text{Fe}/\text{H}] = -0.53$ for SKHB-312,

near the peak in the metallicity distribution of the M31 halo ($[\text{Fe}/\text{H}] = -0.8$; Durrell et al. 2001). Our cluster photometry reaches the main sequence in the F606W (broad V) and F814W (I) bands, allowing a direct estimate of its age. Our analysis is based upon comparisons to photometry of Galactic globular clusters in the same ACS bands, and upon a finely-spaced grid of isochrones calibrated using those clusters.

2. OBSERVATIONS AND DATA REDUCTION

These ACS observations are part of a large program to image the M31 halo, with 3.5 days of exposure time split between the F606W and F814W filters; see Brown et al. (2003) for a description of the data and its reduction. The resulting color-magnitude diagram (CMD) extends more than 1.5 mag below the old main sequence turnoff in the M31 halo. We included the globular cluster SKHB-312 in our field so that the turnoff could also be reached in an M31 cluster (Figure 1a). Using images from the Wide Field Planetary Camera 2 (WFPC2), Holland et al. (1997) determined the SKHB-312 tidal radius to be $\approx 10''$. Although small compared to the ACS field ($\approx 205'' \times 205''$), we placed the cluster near the field edge to minimize the field area contaminated by the cluster in case it appeared more extended in deeper observations. As expected for its high metallicity, SKHB-312 has no known RR Lyrae stars, but it does have 13 long period variables near the RGB tip (Brown et al. 2004); two are within the area we ultimately analyzed.

We measured a more tractable $81'' \times 48''$ subsection around SKHB-312. After constructing a variable point spread function (PSF) from the most isolated stars in the cluster vicinity, we performed aperture and PSF-fitting photometry using the DAOPHOT-II package (Stetson 1987), with zeropoints matching the full field catalog (Brown et al. 2003). Our photometry is in the STMAG system: $m = -2.5 \times \log_{10} f_\lambda - 21.1$ mag.

We constructed our SKHB-312 CMD from stars within an annulus chosen to maximize the signal-to-noise ratio and minimize field contamination. The inner radius (100 pix; 3'') was where the completeness dropped precipitously for stars near the turnoff (Figure 1b); the outer radius (300 pix; 9'') was where the density dropped to twice that in the field (Figure 1c). Because SKHB-312 was near the field edge and the observations were

¹Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS 5-26555. These observations are associated with proposal 9453.

²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; tbrown@stsci.edu, ferguson@stsci.edu, ed-smith@stsci.edu

³Code 681, NASA Goddard Space Flight Center, Greenbelt, MD 20771; randy.a.kimble@nasa.gov, allen.v.sweigart@nasa.gov

⁴European Southern Observatory, Karl-Schwarzschild-Strasse 2, Garching bei München, Germany; arenzini@eso.org

⁵Division of Astronomy, Dpt. of Physics & Astronomy, UCLA, Los Angeles, CA 90095; rmr@astro.ucla.edu

⁶Department of Physics & Astronomy, University of Victoria, P.O. Box 3055, Victoria, BC, V8W 3P6, Canada; davb@uvvm.uvic.ca

dithered, the exposure time was not uniform across the annulus. We discarded the fraction of the annulus ($< 0.5\%$) that had half of the total exposure time, but kept the fraction (14%) that was exposed for 75%; 86% of the annulus was exposed fully.

We used extensive artificial star tests to determine the photometric scatter and completeness as a function of color, luminosity, and field position. Small numbers of stars were repeatedly added to the images with a distribution tracking the cluster light profile (to match these tests to the data), with the noise reflecting the exposure time variations. Only those results within the 100–300 pix annulus were used in our modeling. We then retained those 1720 stars in the SKHB-312 catalog above the 80% completeness magnitude, with the cutoff being a function of position (see Figure 1b), thus producing a much cleaner CMD. For consistency, we treated the cluster and field contamination models the same way (§3).

Our program included ACS observations of five Galactic clusters spanning a wide $[\text{Fe}/\text{H}]$ range, to provide cluster fiducials and to verify the transformation of our isochrones to the ACS bands. The most relevant here are 47 Tuc and NGC5927, bracketing the SKHB-312 metallicity. Our adopted parameters are listed in Table 1; these values give consistency across the Galactic cluster observations and imply one simple transformation of all isochrones to the ACS bands. Note that our isochrones do not include core He diffusion, which would reduce their ages by 10–12% (VandenBerg et al. 2002). After reevaluating the range of parameters in the literature, the Galactic cluster parameters have been updated slightly from those assumed in our earlier papers (Brown 2003; Brown et al. 2003), and the changes have been included in our calibration of the isochrone transformation. The parameters for NGC5927 are highly uncertain, due to its high, spatially variable reddening (Heitsch & Richtler 1999), but the values here are very close to those in Harris (1996). We observed the Galactic clusters with staggered exposure times to increase the dynamic range, but the upper RGB saturated in the shortest possible exposure of 0.5 s.

3. ANALYSIS

We show the CMD of SKHB-312 in Figure 1d. This is the first CMD to resolve the main sequence in a globular cluster of M31. The horizontal branch (HB) consists of a tight red clump (unobscured in Figure 1e), with no extension to a blue HB, while the RGB is very red; these features are distinct from the CMD of the general field population (Brown et al. 2003), and indicate that the cluster is more metal rich than many of the field stars. In this panel, we also plot the ridge lines (*curves*) and HB loci (*points*) of 47 Tuc (*green*) and NGC 5927 (*red*), shifted by the differences in distance and reddening between the clusters and M31 (see Table 1). Instead of applying a constant shift to the entire Galactic cluster fiducial (i.e., its ridge line and HB locus), we applied a correction that more accurately accounts for the ACS response curve. At each point along the fiducial we selected a synthetic spectrum (Castelli & Kurucz 2003), using corresponding points on a theoretical isochrone to determine the appropriate T_{eff} and $\log g$. We then reddened this spectrum with a Fitzpatrick (1999) extinction law, first scaled to the cluster value and then to the M31 value. After folding both through the ACS bands, the difference provided the reddening vector at that point in the fiducial. Accounting for differences in metallicity, the agreement at the HB indicates consistency in the assumed distance moduli, and the relative RGB positions are as expected. However, the subgiant branch of each Galactic cluster is significantly fainter than that of SKHB-312. Inspec-

tion of the CMD and the luminosity function (LF) of each cluster shows that Δm_{F814W} from the HB to the subgiant branch is ≈ 0.2 mag smaller in SKHB-312 than in either of the Galactic clusters; this is an indication that SKHB-312 is 2–3 Gyr younger than those clusters, independent of distance, reddening, and (to a considerable extent) metallicity assumptions.

We used 3 approaches to estimate the age of SKHB-312: fitting by eye of the CMD to calibrated isochrones (as often done in the literature), comparison of the CMD to population models using a Maximum Likelihood statistic, and comparison of the LF to population models using several different statistics. Our models are based upon the isochrones of VandenBerg et al. (2004), which agree well with CMDs of Galactic clusters spanning a wide range of metallicity. These isochrones have been slightly updated from those used in Brown et al. (2003). We also used a new grid of synthetic spectra (Castelli & Kurucz 2003) to calculate the transformation directly from physical parameters (T_{eff} , $\log g$) in the isochrones to magnitudes in the ACS bands (m_{F606W} , m_{F814W}); the updated spectra include changes to the convective overshooting, new opacities, and the availability of alpha-enhanced compositions. As in our earlier work, this transformation includes a small empirical color correction ($\lesssim 0.05$ mag) to force agreement between the transformed isochrones and the ACS CMDs of the Galactic clusters. This correction is smaller and simpler than that used previously (likely due to the improvements in the isochrones and spectral libraries), but the net result is very similar, given the goal is agreement with the same ACS cluster observations. After this correction, the isochrones match the ridge lines within $\lesssim 0.02$ mag until the saturation point in the Galactic clusters. When comparing isochrones to cluster data, we have used isochrones with $[\alpha/\text{Fe}]=0.3$, and transformed them using spectra with the same enhancement. Note that globular clusters tend to show alpha enhancement at all metallicities (Thomas, Maraston, & Bender 2003), compared to the field population.

We show the fitting by eye in Figure 1e, where we compare the SKHB-312 CMD to isochrones of 8, 10, and 12 Gyr. The RGB is very insensitive to age, but sensitive to metallicity. Our transformed isochrones at $[\text{Fe}/\text{H}]=-0.53$ agree well with the RGB of SKHB-312, implying the same metallicity determined by Holland et al. (1997). There is good reason to use the Holland et al. (1997) metallicity: our transformation to the ACS bands cannot be verified on the upper RGB, where the lever on metallicity is greatest, due to the saturation of such stars in the ACS observations of Galactic clusters, while the Holland et al. (1997) work was based upon WFPC2 data, which had a very reliable transformation between ground and space bands (Holtzman et al. 1995). Clearly the main sequence turnoff and

TABLE 1: M31 and Galactic cluster parameters

Name	$(m - M)_V$ (mag)	$E(B - V)$ (mag)	$[\text{Fe}/\text{H}]$	age (Gyr)
SKHB-312	24.68 ^a	0.08 ^b	-0.53 ^c	10.0
47 Tuc	13.27 ^d	0.024 ^e	-0.70 ^f	12.5
NGC5927	15.85	0.42	-0.37 ^g	12.5

^aFreedman & Madore (1990).

^bSchlegel et al. (1998).

^cHolland et al. (1997).


^dZoccali et al. (2001).

^eGratton et al. (2003).

^fKraft & Ivans (2003).

^gPuzia et al. (2002).

FIG. 1— (a) A $24'' \times 24''$ subsection of the F606W image, centered on SKHB-312, with a logarithmic stretch. A star near the turnoff ($m_{F814W} = 29$ mag, $m_{F606W} - m_{F814W} = -0.4$ mag) is circled for reference. (b) The completeness of our photometry as a function of m_{F814W} (labeled) and distance from the cluster. (c) The stellar density as a function of distance from the cluster, for stars above the turnoff ($m_{F814W} \leq 29$), normalized to unity (*dashed*) beyond the tidal radius of the cluster, uncorrected for incompleteness. Our CMD analysis is restricted to the annulus spanning 100 pix – 300 pix (300 pix – 3000 pix), i.e., from the point where the star counts roll over due to severe crowding, to the point where the field contamination is 50%. (d) The CMD from the annulus shown in the previous panel. Note the tight clump of red HB stars (unobscured in the next panel) and the red RGB locus, compared to the CMD of the field population (Brown et al. 2003). The turnoff is clearly detected. The ridge lines (*curves*) and HB locus (*points*) from two Galactic clusters (*labeled*) are overplotted. The clusters bracket the metallicity of SKHB-312, yet the turnoff of each is fainter than that of SKHB-312. The luminosity difference between the HB and the subgiant branch is ≈ 0.2 mag smaller in SKHB-312 than in the Galactic clusters, indicating an age 2–3 Gyr younger. (e) The CMD of SKHB-312 with a 10 Gyr isochrone (*yellow*) at the cluster metallicity shows good agreement; 8 Gyr (*green*) is clearly too young, and 12 Gyr (*red*) is too old. (f) A 10 Gyr isochrone plotted with the same halo contamination, scattering, and completeness (*shaded*) as in the data; beyond the CMD region used for fitting, the SKHB-312 data are shown. This 10 Gyr model agrees well with the same region shown in the previous panels.



subgiant branch in Figure 1e are too bright at 8 Gyr, and too faint at 12 Gyr, implying an age of $\approx 10 \pm 1$ Gyr for SKHB-312. However, fitting by eye ignores two subtle effects that might bias the fit toward younger ages: contamination by field stars, which are known to span a wide age range (Brown et al. 2003), and blends of stars in the crowded photometry, which can brighten the main sequence turnoff and subgiant branch. These effects are included in our quantitative modeling below.

We compared CMD models to the data using a Maximum Likelihood statistic: $\sum \ln P_i$, where P_i is the probability of a star falling at its observed CMD location, given the model under consideration. Each model gives the expected distribution of stars in the CMD, on a 0.02 mag grid, so that the probability of a star being observed in a given bin (P_i) is the fraction of the model falling in that bin (see, e.g., Dolphin 2002). The probability that all of the stars were observed in their given CMD bins is the product of the individual probabilities, and the Maximum Likelihood statistic is a logarithmic score reflecting the probability of observing the SKHB-312 CMD under a given model. We focused on the CMD region highlighted in Figure 1f: $-0.8 \leq (m_{F606W} - m_{M814W}) \leq -0.1$ mag and $26.5 \leq m_{F814W} \leq 30.5$ mag, as in Brown et al. (2003), which includes 1456 of the 1720 stars in the SKHB-312 catalog.

To construct each synthetic CMD, we first populated our isochrones with a Salpeter initial mass function (IMF), and then applied the photometric errors, blends, and incompleteness calculated in the artificial star tests, using the Starfish code (Harris & Zaritsky 2001). We then modeled the halo field contamination by applying the photometric errors, blends, and incompleteness of the SKHB-312 annulus to the best-fit combination of isochrones in Brown et al. (2003), which reproduces the halo data very well. Note that one cannot simply statistically subtract the field data from the cluster data, because they have dramatically different noise and completeness. After normalizing the field model to the area of the SKHB-312 annulus, we found that the halo field contamination in the SKHB-312 CMD was 8% by number. We combined the cluster model and field contamination model to create a synthetic CMD for comparison to the data, with the total number of stars normalized to 1456. This was repeated for isochrone ages of 5–15 Gyr with 0.5 Gyr steps, with the likelihood statistic calculated for each, giving a best-fit of 11 Gyr. To evaluate the goodness of fit, we created Monte Carlo realizations of the best-fit model and evaluated the likelihood statistic. The likelihood values for these samples were higher than for the model tested against the real data, suggesting that the models are imperfect (e.g., because of binaries, blue

stragglers, a non-Salpeter IMF, or remaining imperfections in the isochrones). We reproduced the likelihood values seen for the best fit by distributing 60 stars (4% of the sample) randomly within the CMD region being fit. We used the distribution of the Monte Carlo scores (including 60 randomly distributed stars) to assign confidence intervals to the ages measured from the real data. The results imply age constraints of $9.5 \leq \text{age} \leq 12.5$ Gyr at 68.3% confidence and $9.0 \leq \text{age} \leq 14.0$ Gyr at 99.7% confidence. Thus, the likelihood fitting of the CMD is consistent with our “by eye” age estimate of 10 Gyr. Figure 1f shows the CMD of SKHB-312 outside of the fitting region, and a realization of the 10 Gyr model within the fitting region. Comparison of the model to the data (Figure 1e) shows good agreement.

Finally we fit the SKHB-312 LF, which is often preferred when fitting data with few stars and/or large photometric errors. We collapsed the model CMDs and the data CMD in color, and then compared them using three statistics: χ^2 , χ^2_γ (Mighell 1999), and the likelihood statistic of Dolphin (2002). The best fit in each case, respectively, is 10 Gyr (9–12.5 Gyr at 68.3% confidence), 10.5 Gyr (9–13.0 Gyr at 68.3% confidence), and 10 Gyr (8.5–12.5 Gyr at 68.3% confidence); the confidence intervals come from Monte Carlo tests, comparing the best model against realizations of that model (artificial data), as done in the Maximum Likelihood fitting above. These results are also consistent with the likelihood fitting to the CMD and our fitting by eye, implying an age of ≈ 10 Gyr for SKHB-312.

4. SUMMARY

We have directly determined the age of the M31 cluster SKHB-312, by fitting its deep photometry extending below the main sequence turnoff. These data presented a difficult challenge because the cluster is so compact, and despite our extensive efforts, we do not find tight formal constraints on its age: $10^{+2.5}_{-1}$ Gyr (or ≈ 9 Gyr if core He diffusion were included in the models). However, SKHB-312 appears to be 2–3 Gyr younger than Galactic clusters at similar metallicities. Although these data prove that direct ages can be determined for M31 clusters with existing instrumentation, a program seeking tighter constraints would benefit from the use of the ACS High Resolution Camera and the choice of a less compact cluster.

Support for proposal 9453 was provided by NASA through a grant from STScI, which is operated by AURA, Inc., under NASA contract NAS 5-26555. We are grateful to J. Harris and P. Stetson for providing their codes and assistance.

REFERENCES

- Barmby, P., & Huchra, J.P. 2000, *ApJ*, 531, L29.
 Brown, T.M. 2003, in *The Local Group as an Astrophysical Laboratory*, ed. M. Livio (Cambridge: Cambridge Univ. Press), in press. astro-ph/0308298
 Brown, T.M., Ferguson, H.C., Smith, E., Kimble, R.A., Sweigart, A.V., Renzini, A., & Rich, R.M. 2004, *AJ*, 127, 2738
 Brown, T.M., Ferguson, H.C., Smith, E., Kimble, R.A., Sweigart, A.V., Renzini, A., Rich, R.M., & VandenBerg, D.A. 2003, *ApJ*, 592, L17
 Burstein, D., et al. 2004, *ApJ*, in press. astro-ph/0406564.
 Castelli, F., & Kurucz, R.L. 2003, in *IAU Symposium 210, Modeling of Stellar Atmospheres*, eds. N. Piskunov, W.W. Weiss, & D.F. Gray, poster A20.
 Dolphin, A.E. 2002, *MNRAS*, 332, 91
 Durrell, P.R., Harris, W.E., & Pritchet, C.J. 2001, *ApJ*, 121, 2557
 Fitzpatrick, E.L. 1999, *PASP*, 111, 63
 Freedman, W.L., & Madore, B.F. 1990, *ApJ*, 365, 186
 Gratton, R.G., Bragaglia, A., Carretta, E., Clementini, G., Desidera, S., Grundahl, F., & Lucatello, S. A&A, 408, 529
 Harris, W.E., 1996, *AJ*, 112, 1487
 Harris, J., & Zaritsky, D. 2001, *ApJS*, 136, 25
 Heitsch, E., & Richtler, T. 1999, *A&A*, 347, 455
 Holland, S., Fahlman, G.G., & Richer, H.B. 1997, *AJ*, 114, 1488
 Holtzman, J.A., Burrows, C.J., Casertano, S., Hester, J.J., Trauger, J.T., Watson, A.M., & Worthey, G. 1995, *PASP*, 107, 1065
 Huchra, J.P., Brodie, J.P., & Kent, S.M. 1991, *ApJ*, 370, 495
 Kraft, R.P., & Ivans, I.I. 2003, *PASP*, 115, 143
 Mighell, K.J. 1999, *ApJ*, 518, 380
 Puzia, T.H., Saglia, R.P., Kissler-Patig, M., Maraston, C., Greggio, L., Renzini, A., & Ortolani, S. 2002, *A&A*, 395, 45
 Rich, R.M., Shara, M.M., & Zurek, D. 2001, *AJ*, 122, 842
 Sandage, A.R. 1953, *AJ*, 58, 61
 Sargent, W.L.W., Kowal, C.T., Hartwick, F.D.A., & van den Bergh, S. 1977, *AJ*, 82, 947.
 Schlegel, D.J., Finkbeiner, D.P., & Davis, M. 1998, *ApJ*, 500, 525
 Stetson, P.B. 1987, *PASP*, 99, 191
 Thomas, D., Maraston, C., & Bender, R. 2003, *MNRAS*, 339, 897
 VandenBerg, D.A. 2000, *ApJS*, 129, 315
 VandenBerg, D.A., Bergbusch, P.A., & Dowler, P.D. 2004, in prep.
 VandenBerg, D.A., Richard, O., Michaud, G., & Richer, J. 2002, *ApJ*, 571, 487
 Zoccali, M., et al. 2001, *ApJ*, 553, 733